



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ³ : C07D319/08; A61K 31/365	A1	(11) International Publication Number: WO 83/ 03411 (43) International Publication Date: 13 October 1983 (13.10.83)
<p>(21) International Application Number: PCT/US83/00414</p> <p>(22) International Filing Date: 24 March 1983 (24.03.83)</p> <p>(31) Priority Application Number: PF 3285</p> <p>(32) Priority Date: 24 March 1982 (24.03.82)</p> <p>(33) Priority Country: AU</p> <p>(71) Applicant (<i>for all designated States except US</i>): R.P. SCHERER CORPORATION [US/US]; 2075 West Big Beaver Road, Troy, MI 48099 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (<i>for US only</i>): BATEMAN, Neil, E. [AU/AU]; 23 Edro Avenue, East Brighton, Melbourne, VIC (AU). WOODS, Ross, A. [AU/AU]; Flat 3, 10 Cook Street, West, Brunswick, VIC (AU).</p> <p>(74) Agent: CALLAHAN, James, V.; Allegetti, Newitt, Witcoff & McAndrews, Ltd., 125 South Wacker Drive, Chicago, IL 60606 (US).</p>		<p>(81) Designated States: BR, DK, JP, NO, US.</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: PHARMACEUTICAL COMPOUND AND PROCESS FOR THE MANUFACTURE THEREOF</p> <div data-bbox="785 1768 1444 2107"></div> <p>(57) Abstract</p> <p>An analgesic and anti-inflammatory product comprising a compound having the formula (I), wherein R and R¹ are hydrocarbon radicals and process for the manufacture thereof.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	LI	Liechtenstein
AU	Australia	LK	Sri Lanka
BE	Belgium	LU	Luxembourg
BR	Brazil	MC	Monaco
CF	Central African Republic	MG	Madagascar
CG	Congo	MR	Mauritania
CH	Switzerland	MW	Malawi
CM	Cameroon	NL	Netherlands
DE	Germany, Federal Republic of	NO	Norway
DK	Denmark	RO	Romania
FI	Finland	SE	Sweden
FR	France	SN	Senegal
GA	Gabon	SU	Soviet Union
GB	United Kingdom	TD	Chad
HU	Hungary	TG	Togo
JP	Japan	US	United States of America
KP	Democratic People's Republic of Korea		

-1-

PHARMACEUTICAL COMPOUND AND PROCESS FOR THE
MANUFACTURE THEREOFBACKGROUND OF THE INVENTION - FIELD OF THE
INVENTION AND DESCRIPTION OF THE PRIOR ART

The present invention relates to new pharmaceutical compounds and to a process for the preparation thereof. In particular,
5 the present invention relates to analgesic and anti-inflammatory compounds.

Acetyl salicylic acid, or aspirin as it is more commonly known, is amongst the most widely used of proprietary medicines. Aspirin can be used in the treatment of numerous ailments and is indicated
10 to have analgesic, anti-inflammatory, antipyretic and antirheumatic activity. However, side effects of the drug may limit its application. For example, acetyl salicylic acid may cause gastric irritation and is contra-indicated where such irritation must necessarily be avoided.



-2-

SUMMARY OF THE INVENTION

The object of the present invention is to provide pharmaceutical substances and compositions which exhibit some or all of the advantageous properties of aspirin but in which the
5 contra-indications are minimized.


According to the present invention, it has been discovered that certain esters of acetyl salicylic acid derivatives provide useful pharmaceutical activity.



-3-

DETAILED DESCRIPTION OF
THE PREFERRED EMBODIMENT

The present invention provides a compound of the formula I:




- 5 In formula I, R and R¹ which may be the same or different, represent any suitable radical, such as a hydrocarbon group.

The present invention further provides pharmaceutically acceptable derivatives thereof and bioprecursors therefor. Pharmaceutically acceptable derivatives thereof may include acid
10 addition salts for example.

In a preferred form R may represent an alkyl group. The alkyl group may include from 1 to 15 carbon atoms. The alkyl group may be a straight chain alkyl group. In a preferred form R represents a straight chained alkyl group of from 7 to 11 carbon atoms. In a
15 preferred form of the invention R¹ represents a lower alkyl group, for example, a methyl, ethyl or propyl group. A methyl group has been found to be most satisfactory.

Thus by way of example the compounds of the present invention may be of the formula II:



-4-


where R is a straight chain alkyl radical having from 1 to 15 carbon atoms and preferably from 7 to 11 carbon atoms.

The compounds of the present invention have analgesic and anti-inflammatory activity similar to those of acetyl salicylic acid. It has been found that the following tests provided a valuable guide to the activity of the compounds of the present invention. These tests are the animal model experiments described by L.B. Witkin et al., J. Pharmacol. Exptl. Therap., 133, 400 (1961) and C.A. Winter et al., Proc Soc, Exp. Biol. Med., III, 544 (1962). The use of compounds of the formula I is particularly advantageous since gastric irritation associated with acetyl salicylic acid therapy is avoided or at least minimized.

It is to be understood that this invention encompasses the individual isomeric forms of the compounds as well as mixtures thereof. This includes optically active isomers and racemic mixtures thereof. The benzodioxane derivatives of the present invention may be manufactured by any chemical process known to be useful for the manufacture of chemically analagous compounds.

Compounds of the general formula I may be prepared as a mixture of diastereomers. Such diastereomers are separable by standard techniques e.g. chromatography.

A preferred process for the manufacture of a benzodioxane derivative of the present invention comprises the reaction of an acetyl salicylic acid derivative with a monoglycol ester of the general formula III:



III

wherein R and R^I have the meanings given above. Standard techniques may be utilized in the reaction between acetyl salicylic acid and the monoglycol esters of formula III for the preparation of




-5-

2-alkoxy-4-oxo-1, 3 benzodioxanes as exemplified by C. Ruchardt and S. Rochlitz. Liebigs Ann, Chem., 15 (1974) and G.Y. Paris et al., J. Med. Chem. 23 79 (1980). For example, the acetyl salicylic acid may be activated by conversion to its acylhalide or mixed anhydride to form a 1, 3-benzodioxan-4-one. 1,3-benzodioxan-4-one may be subsequently reacted with monoglycol ester of formula III to produce the compounds of the general formula I. The halide may be selected from a chloride, bromide and iodide. An acid chloride is preferred.


Compounds of the formula III may be prepared via an acylation reaction of a hydroxyketone of the formula IV:



wherein R^1 has the meaning given above; with a carboxylic acid of the formula V:



wherein R has the meaning given above; to yield compounds of the formula VI:




A particularly preferred hydroxyketone is the monohydroxy acetone.

This yields a monoglycol ester of the formula VII:

-6-

This subsequently yields a benzodioxane derivative of the formula II:



Particularly preferred compounds of the present invention which exhibit analgesic and anti-inflammatory activity but with reduced gastric irritation include 2-(1-Decanoyloxyprop-2-oxy)-2-methyl-4-oxo-1,3-benzodioxane, 2-(1-Decanoyloxypropyl-2-oxy)-2-methyl-4-oxo-1,3-benzodioxane, 1-Octanoyloxypropan-2-ol and 2-(1-Octanoyloxyprop-2-oxy)-2-methyl-4-oxo-1,3-benzodioxane.

A suitable acid-addition salt of benzodioxane derivative of the invention is, for example, a salt derived from an organic acid, for example, a hydrochloride, hydrobromide, phosphate or sulphate, or a salt derived from an organic acid, for example an oxalate, lactate, succinate, tartrate, acetate, salicylate, citrate, benzoate, B-naphthoate, or adipate.

The benzodioxane derivatives of the present invention may be administered to animals including man, in the form of a pharmaceutical composition comprising as active ingredient at least one benzodioxane derivative of the present invention, or an acid-addition salt thereof, in association with a pharmaceutically acceptable diluent or carrier therefor. The composition may further comprise a physiologically acceptable excipient, binder, preservative, stabilizer, flavoring or other compounding ingredient. The composition may be prepared in a conventional unit dosage form as called for by accepted pharmaceutical practice. The composition may be included in soft gelatin capsules, two piece hard gelatin shell capsules, tablets, elixirs, suspensions, emulsions or in injectible solutions or suspensions or oily solutions or suspensions.

-7-

The amount of active substances included is selected so as to provide an individual unit dosage, preferably from about 30 milligrams to 1500 milligrams of the active ingredient. Other therapeutically valuable substances may also be included. Caffeine, phenacetin, paracetamol may be used in addition to the active ingredient of the present invention.

The following examples are illustrative of the invention, or of intermediates which may lead to the invention, and represent preferred embodiments but are not to be construed as being limitations thereon. All temperatures are in degrees Celsius.

EXAMPLE 1: 1-Decanoyloxypropan-2-one

Monohydroxyacetone was distilled prior to use. Chloroform was distilled from phosphorous pentoxide prior to use. Decanoyl chloride (65 cm^3 , 0.3146 mole) was added dropwise to stirred solution of monohydroxyacetone (21.8 cm^3 , 0.3146 mole) and anhydrous pyridine (25.4 cm^3 , 0.3146 mole) in anhydrous chloroform (500 cm^3). During the addition the solution was maintained in an ice bath. The mixture was then stirred at room temperature overnight. The solution was washed with water ($4 \times 200 \text{ cm}^3$) and saturated sodium chloride solution (200 cm^3). The organic phase was dried over sodium sulphate and evaporated in vacuo. The resulting oil was distilled to yield 1-decanoyloxypropan-2-one as a mobile liquid, bp $154-60^\circ$ (1mm) (52.4 g, 0.230 mole, 73%), ir (neat) 2860, 1700, 1400, 1340, 1150, 1110 and 1050 cm^{-1} ; pmr (CDCl_3) 4.6 (s, 2 H), 2.4 (t, 2 H), 2.1 (s, 3 H), 1.8-1.0 (bm, 14 H) and 0.9 (t, 3 H) ppm.

EXAMPLE 2: 1-Decanoyloxypropan-2-ol

1-Decanoyloxypropan-2-one (44.3 g, 0.194 mole) was dissolved in a solution of tetrahydrofuran (1100 cm^3) and benzene (200 cm^3). After cooling to 5° , ice water (80 cm^3) was added. Sodium borohydride (11 g, 0.291 mole) was added to the stirred solution in small proportions to maintain the temperature at 5° . After addition the reaction was stirred at 5° for 45 minutes and glacial acetic acid (14 cm^3) was added dropwise. Stirring at 5° was continued for a

-8-

further 30 minutes. Diethyl ether and chloroform (200 cm³ each) were added and the mixture washed with water (2 x 200 cm³), a 1% sodium bicarbonate solution (200 cm³) and brine (200 cm³). The organic phase was dried over sodium sulphate and evaporated to yield 1-decanoyloxypropan-2-ol as a colorless liquid (44.0 g, 0.191 mole, 98%), ir(neat) 3600-3100, 2920, 2860, 1730, 1460, 1375, 1240, 1175, 1110, 1055 cm⁻¹; pmr (CDCl₃) 4.0 (m, 2 H), 3.6 (m, 2 HO (D₂O m, 1 H), 2.3 (t, 2 H), 2.0-1.0 (bm, 17 H) and 0.85 (t, 3 H) ppm.

EXAMPLE 3: 2-(1-Decanoyloxyprop-2-oxy)-2-methyl-4-oxo-1, 3-benzodioxane

Trifluoroacetic anhydride (3.73 cm³, 0.0263 cm³) was added to a suspension of 0-acetylsalicylic acid (4.3 g, 0.239 mole) in anhydrous toluene (100 cm³) at 80°. The solution was stirred at room temperature for 10 minutes. 1-Decanoyloxypropan-2-ol (5 g, 0.0217 mole) in toluene (25 cm³) was added with rapid stirring. The solution was stirred at room temperature for 5 minutes and in ice for 10 minutes. Pyridine (7 cm³, 0.0865 mole) was then added. Additional toluene (100 cm³) was added and the solution extracted with water (2 x 150 cm³), a saturated sodium bicarbonate solution (3 x 150 cm³), dried over sodium sulphate and evaporated. The product was chromatographed on silica gel (600 g) in petroleum ether : ether (80:20) and treated with charcoal in acetone to yield 2-(1-decanoyloxyprop-2-oxy)-2-methyl-4-oxo-1, 3-benzodioxane as a mixture of diastereomers (6.3 g, 0.0161 mole, 67%); ir (neat) 2920, 2850, 1730, 1600, 1580, 1470, 1380, 1310, 1275, 1250, 1150, 1080, 1015, 920 and 760 cm⁻¹; pmr (CDCl₃) 7.9 (dd, 2 H), 7.6-6.8 (m, 3 H), 4.6-3.6 (m, 3H), 2.2 (m, 2 H), 1.85 (s, 3 H), 1.6-1.0 (bm, 17 H) and 0.8 (m, 3 H) ppm.

Anal. calcd. for C₂₂H₃₂O₆ : C, 67.32; H, 8.22%

Found: C, 67.64; H, 8.62%

EXAMPLE 4: 2-(1-Decanoyloxypropyl-2-oxy)-2-methyl-4-oxo-1, 3-benzodioxane

A mixture of 1 equivalent of 0-acetylsalicyloyl chloride, 1-decanoyloxypropan-2-ol and anhydrous pyridine in anhydrous,



-9-

ethanol free, chloroform was heated at reflux for 24 hours. The solution was washed with water, a 1% sodium bicarbonate solution and evaporated. The product was purified as in Example 3 to yield 2-(1-decanoyloxyprop-2-oxy)-2-methyl-4-oxo-1, 3-benzodioxane identical to that in Example 3.

EXAMPLE 5: 1-Octanoyloxypropan-2-ol

By replacing decanoyl chloride in Example with octanoyl chloride, 1-octanoyoxypropan-1-one and subsequently 1-octanoyloxypropan-2-ol, as in Example 2, can be obtained.

10 EXAMPLE 6: 2-(1-Octanoyloxyprop-2-oxy)-2-methyl-4-oxo-1, 3-benzodioxane


By replacing 1-décanoyloxypropan-2-ol in Example 3 with 1-octanoyloxypropan-2-ol, 2-(1-octanoyloxyprop-2-oxy)-2-methyl-4-oxo-1, 3-benzodioxane can be obtained.

15 Finally, it is to be understood that various other modifications and/or alterations may be made without departing from the spirit of the present invention as outlined herein.



CLAIMS:

1. An analgesic and anti-inflammatory compound having the formula:



wherein R and R¹ are hydrocarbon radicals.

2. The product of Claim 1 wherein R is an alkyl group having 1 to 15 carbon atoms.

3. The product of Claim 2 wherein said alkyl group is a straight chain alkyl group.

4. The product of Claim 3 wherein said straight chain alkyl group has 7 to 11 carbon atoms.

5. The product of Claim 1 wherein R¹ is a lower alkyl group.

6. The product of Claim 5 wherein R¹ is a methyl, ethyl, or propyl group.

7. A process for preparing an analgesic and anti-inflammatory compound comprising the step of reacting an acetyl salicylic acid derivative with a monoglycol ester having the formula:



- 5 wherein R and R¹ are hydrocarbon radicals.



INTERNATIONAL SEARCH REPORT

International Application No PCT/US83/00414

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *

According to International Patent Classification (IPC) or to both National Classification and IPC

Int. Cl 3 C07D 319/08 US Cl. 549/274
A61K 31/365

II. FIELDS SEARCHED

Minimum Documentation Searched *

Classification System

Classification Symbols

US

549/274

560/66

260/410.5

424/230

Documentation Searched other than Minimum Documentation
to the extent that such Documents are included in the Fields Searched *

Chemical Abstracts, Volumes 1-97- 1,2-Propanediol, Esters;
Glycols, Esters; Antipyretics; Analgesics; Chemical
Abstracts-On-Line

III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴

Category *	Citation of Document, ¹⁴ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
Y	US,A, 2,070,240, published 9 February 1937 Ruben	1-6
Y	US,A, 2,151,185, published 21 March 1939 Carruthers	1-6
Y	US,A, 3,012,064, published 5 December 1961 Hertling	1-6
Y	US,A, 3,644,424, published 22 February 1972 Sherlock	1-6
Y	US,A, 3,652,665, published 28 March 1972 Shen	1-6
Y	US,A, 3,988,446, published 26 October 1976 Paris	1-6
Y	US,A, 4,046,887, published 6 September 1977 Paris	1-7
Y	Journal of Medicinal Chemistry, Vol.23, No. 1, issued 1980 (Easton, Pennsylvania), Gerard Y. Paris et al., "Glycerides as Prodrugs. 2. 1,3-dialkanoyl-2-(2-methyl- 4-oxo-1,3-benzodioxan-2-yl)glycerides (Cyclic Aspirin Triglycerides) as Antiinflammatory Agents", see pp. 79-82.	1-7

* Special categories of cited documents: ¹⁵

"A" document defining the general state of the art which is not
considered to be of particular relevance

"E" earlier document but published on or after the international
filing date

"L" document which may throw doubts on priority claim(s) or
which is cited to establish the publication date of another
citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or
other means

"P" document published prior to the international filing date but
later than the priority date claimed

"T" later document published after the international filing date
or priority date and not in conflict with the application but
cited to understand the principle or theory underlying the
invention

"X" document of particular relevance; the claimed invention
cannot be considered novel or cannot be considered to
involve an inventive step

"Y" document of particular relevance; the claimed invention
cannot be considered to involve an inventive step when the
document is combined with one or more other such docu-
ments, such combination being obvious to a person skilled
in the art.

"&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search *

06 June 1983

Date of Mailing of this International Search Report *

18 JUL 1983

International Searching Authority *

RO/US

Signature of Authorized Officer ¹⁹

Norma S. Milestone

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, ¹⁶ with Indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No ¹⁸
------------	--	------------------------------------

A	Justus Liebigs Annalen der Chemie, issued 1974 (West Germany), Christoph Ruchardt et al., "Zur ambivalenten Reaktivitat des O-Acetylsalicylsaurechlorids", see pp.15-23.	1-7
---	--	-----

A	The Journal of Pharmacology and Experimental Therapeutics, Vol. 133, issued 1961 (Baltimore, Maryland), L.B. Witkl. "Pharmacology of 2-Amino-indane Hydrochloride (SU-8629): A Potent Non-narcotic Analgesic", see pp. 400-408.	1-6
---	---	-----

A	Proceedings of the Society for Experimental Biology and Medicine, Vol. III, No. 3, issued December 1962 (Utica, New York), Charles A. Winter et al., "Carrageenin-Induced Edema in Hind Paw of the Rat as an Assay for Antiinflammatory Drugs", see pp. 544-547.	1-6
---	--	-----